



IAAF BETA-2 AGONISTS PROTOCOL

In response to a generally recognised increase in the use of beta-2 agonists by athletes, the IAAF Council has endorsed a recommendation from the IAAF Medical and Anti-Doping Commission to require more detailed information from applicants who seek an exemption for the use of beta-2 agonists for asthma and/or exercise-induced asthma or exercise-induced bronchoconstriction, and/or other clinical respiratory diseases requiring this specific treatment.

Applicants seeking the use of the four beta-2 agonists (formeterol, salbutamol, salmeterol and terbutaline) will be required to submit the following information:

- (i) An IAAF TUE Abbreviated Application Form including detailed medical records. See www.iaaf.org/antidoping to download a copy of the TUE form.
- (ii) Objective evidence of asthma and/or exercise-induced asthma or exercise-induced bronchoconstriction and/or other clinical respiratory diseases requiring this specific treatment through the provision of test results and supporting documentation, as set out below (see Provocation Test Results). All tests should have been performed since 1 January 2005.

Note: all TUE Abbreviated Application Forms and supporting documentation must be submitted to the IAAF in either English or French.

1. Detailed Medical Records

The TUE Abbreviated Application Form needs accompanying Medical information and records, including (as suggested by the enclosed form):

- A precise diagnosis of the individual's condition requiring the use of beta-2 agonists.
 - all relevant information regarding the individual concerned and his condition:
 - age of onset
 - symptoms suggesting airway obstruction following exercise, upper respiratory infection at rest and at night and/or during the pollen season
 - identified triggering factors
 - past history of atopic disorders and/or childhood asthma
 - past physical examinations
 - results of skin prick tests or RAST to document the presence of allergic hypersensitivity

- Any specific information concerning the individual's coughing during or postexercise, dyspnoea, shortness of breath, wheezing, chest tightness or excess sputum.
- Details of all consultations with physicians qualified in the treatment of asthma and details of any attendance in hospital emergency departments for treatment or admission to hospital for treatment of acute exacerbation of asthma.
- Details of the individual's currently prescribed medication and any other medication prescribed in the last 6 months. Details of medication in the 3 months prior to provocation tests (see below) must also be notified.

2. Provocation Test Results

In addition to the detailed medical records, athletes must also present a positive test result from one of the following recognised provocation tests:

a) Bronchodilator test:

A positive test result shall be defined as:

- a 12 % or greater increase in FEV₁ calculated as a percentage of the baseline FEV₁ after the administration of an inhaled permitted beta-2 agonist.

Submission of worksheets and graphic evidence (spirometry or flow volume tracings) is useful.

b) Bronchial provocation test:

A Bronchial provocation test will take the form of an exercise challenge in the laboratory or an exercise test in the field or a eucapnic voluntary hyperpnea test in the laboratory (EVH).

A positive test result will be obtained if airway hyper-responsiveness (AHR) is confirmed with a fall of 10% or more in FEV₁ within 30 minutes of finishing the challenge.

Submission of worksheets and graphic evidence (spirometry or flow volume tracings) is useful.

c) Bronchial provocation test with inhaled methacholine:

A positive test result will be obtained if AHR is confirmed with:

c1 When the subject is not taking inhaled glucocorticosteroids:

- a PC₂₀ FEV₁ equal to or less than 4 mg/ml;

OR

- a PD₂₀ FEV₁ equal to or less than a cumulative dose of 2 micromol or 400 micrograms or 40 breath units in steroid-naïve subjects.

c2 In the case of individuals on daily inhaled corticosteroid treatment of more than 3 months duration:

- a PC₂₀ FEV₁ equal to or less than 13.2 mg/ml;

OR

- a PD₂₀ FEV₁ equal to or less than a cumulative dose of 6.6 micromol, or equal to or less than 1320 micrograms or 130 breath units.

Submission of:

- **Details of medication in the 3 months prior to provocation tests (see below) must also be notified.**
- **The physician shall provide exact calculated value of PC₂₀ or PD₂₀ which is mandatory.**
- **Submission of worksheets and graphic evidence (spirometry or flow volume tracings) is useful.**

d) Bronchoconstrictor test:

A positive test result for a bronchoconstriction test is defined as a fall of 15 % or more in FEV₁ after the subject inhaling a hypertonic aerosol (4.5% saline commonly used).

Submission of worksheets and graphic evidence (spirometry or flow volume tracings) is useful.

Note: Peak Expiratory Flow Rate (PEFR) measurements will not be accepted. Submission of bronchial provocation tests using pharmacological agents other than methacholine (e.g. carbachol, histamine or adenosine monophosphate) will not be accepted.

Recommendations prior to the challenge test

To provide the optimal test conditions to achieve a positive response, it is recommended that some medications be withheld for a period of 8-96 hours before the bronchial provocation test, as indicated below:

- No short-acting bronchodilators, sodium chromoglycate, nedocromil sodium, ipratropium bromide or oxitropium or tiotropium bromide for 8 hours.
- No long-acting bronchodilators or antihistamines for 48 hours.
- No leukotriene antagonists for 4 days.
- Inhaled steroids should not be administered on the day of the test.
- No caffeine should be taken on the morning of the test.
- No vigorous exercise to be taken for at least 4 hours prior to the test and preferably none on the day of test at all.

Well-controlled asthma with negative response to all the tests

In the exceptional case of an athlete with known, but well-controlled, asthma reporting “negative” results from the bronchial provocation tests, while applying for the use of inhaled beta2-agonists, the following documentation must be enclosed, in addition to negative results obtained in the bronchial provocation tests: a complete report from personal physician on medical history, including reports on hospital emergency department attendance or admission for acute exacerbations of asthma and on previous treatment with different medications, including systemic corticosteroids, and any additional information that might be helpful for the diagnosis.

Abbreviations

FEV₁ = Forced expired volume in one second.

PC₂₀ FEV₁ = Is the provocative concentration of methacholine causing a 20% fall in FEV₁

PD₂₀ FEV₁ = Is the provocative dose of methacholine causing a 20% fall in FEV₁

References:

1. Anderson SD. Pulmonary function testing and bronchial provocation. In: Altman LC (ed) Allergy in Primary Care. W.B. Saunders Ltd, 2000: 49-64.
2. Anderson D, Argyros G J, Magnussen H, Holzer K. Provocation by eucapnic voluntary hyperpnoea to identify exercise induced bronchoconstriction. Br J Sports Med 2001; 35: 344-347.
3. Anderson SD, Brannan JD. Methods for 'indirect' challenge tests including exercise, eucapnic voluntary hyperpnea and hypertonic aerosols. Clin Rev Allergy Immunol 2003; 24: 63-90.
4. Cockcroft DW, Davis BE, Todd DC, Smycniuk AJ. Methacholine challenge: >comparison of two methods. Chest 2005; 125:839-844
5. Crapo RO, Casaburi R, Coates AL, et al. Guidelines for methacoline and exercise challenge testing-1999. Am J Respir Crt Care Med, 2000; 161:309-29.
6. Gold WM. Pulmonary function testing. IN. Murray JF, Nadel JA, (eds.) Textbook of Respiratory Medicine. 2nd ed. WB Saunders Co, Philadelphia 1994: 798-893.
7. Medical Commission. International Olympic Committee. Beta2 adrenoceptor agonists and the Olympic Winter Games in Salt Lake City.
8. Medical Commission. International Olympic Committee. Beta2 adrenoceptor agonists and the Olympic Games in Turin 2006. http://multimedia.olympic.org/pdf/en_report_981.pdf Accessed on 15th December 2005.
9. Quanjer PH, Tammeling JE, Cotes OF, Pedersen R, Peslin R, Yemault JC. Lung volumes and forced ventilatory flows. Eur Respir J 1993; 6:5-40.
10. Rundell KW, Anderson SD, Spiering BA, Judelson DA. Field exercise vs laboratory eucapnic voluntary hyperventilation to identify airway hyperresponsiveness in elite cold weather athletes. Chest 2004; 125:909-15.
11. Smith CM, Anderson SD. Inhalational challenge using hypertonic saline in asthmatic subjects: a comparison with responses to hyperpnoea, methacholine and water. Eur Respir J 1990; 3: 144-51.
12. Sterk PJ, Fabbri LM, Quanjer PH, et al. Airway responsiveness: Standardized challenge testing with pharmacological, physical and sensitizing stimuli in adults. Eur Respir J, 1993 ; 6 (Suppl 16):53-83.